


Amendments to the Claims:

This listing of claims will replace all prior versions and listings of claims in the application:

Listing of Claims:

Claims 1-2 (canceled).


3. (currently amended) A compound of Claim 2 having the formula I




or a pharmaceutically acceptable salt thereof.

wherein:

R^1 and R^2 are independently selected from the group consisting of H, $(CH_2)_nAr$, COR^4 , $(CH_2)_n$ heteroaryl, $(CH_2)_n$ heterocyclyl, C_1 - C_{10} alkyl, C_3 - C_{10} cycloalkyl, C_2 - C_{10} alkenyl, and C_2 - C_{10} alkynyl, wherein n is 0, 1, 2, or 3, and the $(CH_2)_nAr$, $(CH_2)_n$ heteroaryl, alkyl, cycloalkyl, alkenyl, and alkynyl groups are optionally substituted by up to 5 groups selected from NR^4R^5 , $N^+(O)R^4R^5$, $N^+R^4R^5R^6Y^-$, alkyl, phenyl, substituted phenyl, $(CH_2)_n$ heteroaryl, hydroxy, alkoxy, phenoxy, thiol, thioalkyl, halo, COR^4 , CO_2R^4 , $CONR^4R^5$, $SO_2NR^4R^5$, SO_3R^4 , PO_3R^4 , aldehyde, nitrile, nitro, heteroaryloxy, $T(CH_2)_mQR^4$,



$T(CH_2)_mC(CH_2)_mQR^4$,



$C(O)T(CH_2)_mQR^4$, $NHC(O)T(CH_2)_mQR^4$, $T(CH_2)_mC(O)NR^4NR^5$, or $T(CH_2)_mCO_2R^4$ wherein each m is independently 1-6, T is O, S, NR^4 , $N^+(O)R^4$, $N^+R^4R^6Y^-$, or CR^4R^5 , and Q is O, S, NR^5 , $N^+(O)R^5$, or $N^+R^5R^6Y^-$;

R^4 and R^5 are each independently selected from the group consisting of hydrogen, C_1 - C_6 alkyl, substituted alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, $N(C_1$ - C_6 alkyl)_{1 or 2}, $(CH_2)_nAr$,

C₃-C₁₀ cycloalkyl, heterocyclyl, and heteroaryl, or R⁴ and R⁵ together with the nitrogen to which they are attached optionally form a ring having 3 to 7 carbon atoms and said ring optionally contains 1, 2, or 3 heteroatoms selected from the group consisting of nitrogen, substituted nitrogen, oxygen, and sulfur;

when R⁴ and R⁵ together with the nitrogen to which they are attached form a ring, the said ring is optionally substituted by 1 to 3 groups selected from OH, OR⁴, NR⁴R⁵, (CH₂)_mOR⁴,

(CH₂)_mNR⁴R⁵, T-(CH₂)_mQR⁴, CO-T-(CH₂)_mQR⁴, NH(CO)T(CH₂)_mQR⁴, T-(CH₂)_mCO₂R⁴, or T(CH₂)_mCONR⁴R⁵;

R⁶ is alkyl; and

Y is a halo counter-ion.

4. (currently amended) [[A]] The compound or pharmaceutically acceptable salt thereof of Claim 3 wherein R¹ is phenyl or substituted phenyl, pyridyl or substituted pyridyl.

5. (currently amended) [[A]] The compound or pharmaceutically acceptable salt thereof of Claim 4 wherein R² is an alkyl, substituted alkyl, or cycloalkyl unsubstituted or substituted.

6. (original) A compound selected from:

1-Methyl-7-[4-(pyrazol-1-yl)phenylamino]pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

1-Methyl-7-[4-(4-methylpiperazin-1-yl)phenylamino]pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

1-Methyl-7-[4-(4-hydroxypiperidin-1-yl)phenylamino]pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

1-Methyl-7-[4-[4-(dimethylamino)piperidin-1-yl]phenylamino]pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

1-Isopropyl-7-[4-(pyrazol-1-yl)phenylamino]pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

1-Isopropyl-7-[4-(4-methylpiperazin-1-yl)phenylamino]pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

1-Isopropyl-7-[4-(4-hydroxypiperidin-1-yl)phenylamino]pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

1-Isopropyl-7-[4-[4-(dimethylamino)piperidin-1-yl]phenylamino]pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

1-Bicyclo[2.2.1]hept-2-yl-7-[4-(pyrazol-1-yl)phenylamino]pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one (exo);

1-Bicyclo[2.2.1]hept-2-yl-7-[4-(4-methylpiperazin-1-yl)phenylamino]pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one (exo);

1-Bicyclo[2.2.1]hept-2-yl-7-[4-(4-hydroxypiperidin-1-yl)phenylamino]pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one (exo);

1-Bicyclo[2.2.1]hept-2-yl-7-[4-[4-(dimethylamino)piperidin-1-yl]phenylamino]pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one (exo);

7-[4-(4-Aminoacetyl-piperazin-1-yl)-phenylamino]-1-cyclopentyl-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
 7-{4-[4-(2-Amino-4-methyl-pentanoyl)-piperazin-1-yl]-phenylamino}-1-cyclopentyl-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
 1-Methyl-7-{4-[4-(3-morpholin-4-ylpropyl)piperidin-1-yl]phenylamino}pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
 1-Isopropyl-7-{4-[4-(3-morpholin-4-ylpropyl)piperidin-1-yl]phenylamino}pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
 1-Cyclopentyl-7-{4-[4-(3-morpholin-4-ylpropyl)piperidin-1-yl]phenylamino}pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
 1-Bicyclo[2.2.1]hept-2-yl-7-{4-[4-(3-morpholin-4-ylpropyl)piperidin-1-yl]phenylamino}pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one (exo);
 1-Cyclopentyl-7-(4-methanesulfonyl-phenylamino)-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
 1-Cyclopentyl-7-(4-fluoro-3-methyl-phenylamino)-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
 7-[4-(3-Amino-pyrrolidin-1-yl)-phenylamino]-1-cyclopentyl-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
 1-Cyclopentyl-7-(4-piperazin-1-yl-phenylamino)-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
 1-Cyclopentyl-7-[4-(5-methyl-hexahydro-pyrrolo[3,4-*c*]pyrrol-2-yl)-phenylamino]-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
 7-[4-(4-Acetyl-piperazin-1-yl)-phenylamino]-1-cycloheptyl-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one; and
 1-Cyclopentyl-7-(pyridin-4-ylamino)pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one.

Claims 7-8 (canceled).

9. (original) A compound selected from:

1-Methyl-7-[4-(pyrazol-1-yl)phenylamino]-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
 1-Methyl-7-[4-(4-methylpiperazin-1-yl)phenylamino]-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
 1-Methyl-7-[4-(4-hydroxypiperidin-1-yl)phenylamino]-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
 1-Methyl-7-{4-[4-(dimethylamino)piperidin-1-yl]phenylamino}-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
 1-Isopropyl-7-[4-(pyrazol-1-yl)phenylamino]-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
 1-Isopropyl-7-[4-(4-methylpiperazin-1-yl)phenylamino]-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
 1-Isopropyl-7-[4-(4-hydroxypiperidin-1-yl)phenylamino]-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

1-Isopropyl-7-{4-[4-(dimethylamino)piperidin-1-yl]phenylamino}-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

1-Bicyclo[2.2.1]hept-2-yl-7-[4-(pyrazol-1-yl)phenylamino]-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one (exo);

1-Bicyclo[2.2.1]hept-2-yl-7-[4-(4-methylpiperazin-1-yl)phenylamino]-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one (exo);

1-Bicyclo[2.2.1]hept-2-yl-7-[4-(4-hydroxypiperidin-1-yl)phenylamino]-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one (exo);

1-Bicyclo[2.2.1]hept-2-yl-7-[4-[4-(dimethylamino)piperidin-1-yl]phenylamino]-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one (exo);

7-[4-(4-Aminoacetyl-piperazin-1-yl)-phenylamino]-1-cyclopentyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

7-{4-[4-(2-Amino-4-methyl-pentanoyl)-piperazin-1-yl]-phenylamino}-1-cyclopentyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

1-Methyl-7-{4-[4-(3-morpholin-4-ylpropyl)piperidin-1-yl]phenylamino}-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

1-Isopropyl-7-{4-[4-(3-morpholin-4-ylpropyl)piperidin-1-yl]phenylamino}-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

1-Cyclopentyl-7-{4-[4-(3-morpholin-4-ylpropyl)piperidin-1-yl]phenylamino}-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

1-Bicyclo[2.2.1]hept-2-yl-7-[4-[4-(3-morpholin-4-ylpropyl)piperidin-1-yl]phenylamino]-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one (exo);

1-Cyclopentyl-7-(pyridin-4-ylamino)-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

1-Cyclopentyl-7-(4-methanesulfonyl-phenylamino)-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

1-Cyclopentyl-7-(4-fluoro-3-methyl-phenylamino)-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

7-[4-(3-Amino-pyrrolidin-1-yl)-phenylamino]-1-cyclopentyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

7-[4-(4-Acetyl-piperazin-1-yl)-phenylamino]-1-cyclopentyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

1-Cyclopentyl-7-(4-piperazin-1-yl-phenylamino)-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

1-Cyclopentyl-7-[4-(5-methyl-hexahydro-pyrrolo[3,4-*c*]pyrrol-2-yl)-phenylamino]-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

7-[4-(4-Aminoacetyl-piperazin-1-yl)-phenylamino]-3-(3,5-dimethoxy-phenyl)-1-ethyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

7-[4-(4-Aminoacetyl-piperazin-1-yl)-phenylamino]-3-(2-chloro-3,5-dimethoxy-phenyl)-1-ethyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

7-[4-(4-Aminoacetyl-piperazin-1-yl)-phenylamino]-3-(2,6-dichloro-3,5-dimethoxy-phenyl)-1-ethyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

7-[4-(4-Aminoacetyl-piperazin-1-yl)-phenylamino]-3-(2-methyl-3,5-dimethoxy-phenyl)-1-ethyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

7-[4-(4-Aminoacetyl-piperazin-1-yl)-phenylamino]-3-(2,6-dimethyl-3,5-dimethoxy-phenyl)-1-ethyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

7-[4-(2-Diethylamino-ethoxy)-phenylamino]-3-(3,5-dimethoxy-phenyl)-1-ethyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

7-[4-(2-Diethylamino-ethoxy)-phenylamino]-3-(2-chloro-3,5-dimethoxy-phenyl)-1-ethyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

7-[4-(2-Diethylamino-ethoxy)-phenylamino]-3-(2,6-dichloro-3,5-dimethoxy-phenyl)-1-ethyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

7-[4-(2-Diethylamino-ethoxy)-phenylamino]-3-(2-methyl-3,5-dimethoxy-phenyl)-1-ethyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

7-[4-(2-Diethylamino-ethoxy)-phenylamino]-3-(2,6-dimethyl-3,5-dimethoxy-phenyl)-1-ethyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

7-(4-Diethylamino-butylamino)-3-(3,5-dimethoxy-phenyl)-1-ethyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

7-(4-Diethylamino-butylamino)-3-(2-chloro-3,5-dimethoxy-phenyl)-1-ethyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

7-(4-Diethylamino-butylamino)-3-(2,6-dichloro-3,5-dimethoxy-phenyl)-1-ethyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

7-(4-Diethylamino-butylamino)-3-(2-methyl-3,5-dimethoxy-phenyl)-1-ethyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

7-(4-Diethylamino-butylamino)-3-(2,6-dimethyl-3,5-dimethoxy-phenyl)-1-ethyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

7-(Pyridin-4-ylamino)-3-(3,5-dimethoxy-phenyl)-1-ethyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

7-(Pyridin-4-ylamino)-3-(2-chloro-3,5-dimethoxy-phenyl)-1-ethyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;


7-(Pyridin-4-ylamino)-3-(2,6-dichloro-3,5-dimethoxy-phenyl)-1-ethyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

7-(Pyridin-4-ylamino)-3-(2,6-dimethyl-3,5-dimethoxy-phenyl)-1-ethyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

7-(Pyridin-4-ylamino)-3-(2-methyl-3,5-dimethoxy-phenyl)-1-ethyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

7-(Pyridin-4-ylamino)-3-(2,6-dichloro-3,5-dimethoxy-phenyl)-1-cyclopentyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
 3-(2-Chloro-3,5-dimethoxy-phenyl)-7-(4-diethylamino-butylamino)-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
 3-(2-Chloro-3,5-dimethoxy-phenyl)-7-[4-(2-diethylamino-ethoxy)-phenylamino]-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
 3-(2-Chloro-3,5-dimethoxy-phenyl)-7-(pyridin-4-ylamino)-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
 3-(3,5-Dimethoxy-phenyl)-7-(pyridin-4-ylamino)-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
 7-[4-(2-Diethylamino-ethoxy)-phenylamino]-3-(3,5-dimethoxy-phenyl)-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
 3-(2,6-Dichloro-3,5-dimethoxy-phenyl)-7-(pyridin-4-ylamino)-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one; and
 3-(2,6-Dichloro-3,5-dimethoxy-phenyl)-7-[4-(2-diethylamino-ethoxy)-phenylamino]-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one.


10. (currently amended) A compound of ~~Claim 2 having the~~ formula II




or a pharmaceutically acceptable salt thereof,

wherein:


R¹ is selected from the group consisting of H, (CH₂)_nAr, COR⁴, (CH₂)_nheteroaryl, (CH₂)_nheterocyclyl, C₁-C₁₀ alkyl, C₃-C₁₀ cycloalkyl, C₂-C₁₀ alkenyl, and C₂-C₁₀ alkynyl, wherein n is 0, 1, 2, or 3, and the (CH₂)_nAr, (CH₂)_nheteroaryl, alkyl, cycloalkyl, alkenyl, and alkynyl groups are optionally substituted by up to 5 groups selected from NR⁴R⁵, N⁺(O)R⁴R⁵, N⁺R⁴R⁵R⁶Y⁻, alkyl, phenyl, substituted phenyl, (CH₂)_nheteroaryl, hydroxy, alkoxy, phenoxy, thiol, thioalkyl, halo, COR⁴, CO₂R⁴, CONR⁴R⁵, SO₂NR⁴R⁵, SO₃R⁴, PO₃R⁴, aldehyde, nitrile, nitro, heteroaryloxy, T(CH₂)_mQR⁴,





$\text{C(O)T(CH}_2\text{)}_m\text{QR}^4$, $\text{NHC(O)T(CH}_2\text{)}_m\text{QR}^4$, $\text{T(CH}_2\text{)}_m\text{C(O)NR}^4\text{NR}^5$, or $\text{T(CH}_2\text{)}_m\text{CO}_2\text{R}^4$ wherein each m is independently 1-6, T is O , S , NR^4 , $\text{N}^+(\text{O})\text{R}^4$, $\text{N}^+\text{R}^4\text{R}^6\text{Y}^-$, or CR^4R^5 , and Q is O , S , NR^5 , $\text{N}^+(\text{O})\text{R}^5$, or $\text{N}^+\text{R}^5\text{R}^6\text{Y}^-$;

R^3 has the meanings of R^1 , wherein R^1 is as defined above, as well as OH , NR^4R^5 , COOR^4 , OR^4 , CONR^4R^5 , $\text{SO}_2\text{NR}^4\text{R}^5$, SO_3R^4 , PO_3R^4 , $\text{T(CH}_2\text{)}_m\text{QR}^4$,



wherein T and Q are as defined above;

R^4 and R^5 are each independently selected from the group consisting of hydrogen, $\text{C}_1\text{-C}_6$ alkyl, substituted alkyl, $\text{C}_2\text{-C}_6$ alkenyl, $\text{C}_2\text{-C}_6$ alkynyl, $\text{N(C}_1\text{-C}_6\text{alkyl)}_1$ or 2 , $(\text{CH}_2)_n\text{Ar}$,

$\text{C}_3\text{-C}_{10}$ cycloalkyl, heterocyclyl, and heteroaryl, or R^4 and R^5 together with the nitrogen to which they are attached optionally form a ring having 3 to 7 carbon atoms and said ring optionally contains 1, 2, or 3 heteroatoms selected from the group consisting of nitrogen, substituted nitrogen, oxygen, and sulfur;

when R^4 and R^5 together with the nitrogen to which they are attached form a ring, the said ring is optionally substituted by 1 to 3 groups selected from OH , OR^4 , NR^4R^5 , $(\text{CH}_2)_m\text{OR}^4$,

$(\text{CH}_2)_m\text{NR}^4\text{R}^5$, $\text{T-(CH}_2\text{)}_m\text{QR}^4$, $\text{CO-T-(CH}_2\text{)}_m\text{QR}^4$, $\text{NH(CO)T(CH}_2\text{)}_m\text{QR}^4$, $\text{T-(CH}_2\text{)}_m\text{CO}_2\text{R}^4$, or $\text{T(CH}_2\text{)}_m\text{CONR}^4\text{R}^5$;

R^6 is alkyl; and

Y is a halo counter-ion.

11. (original) A compound selected from:


1-[7-[4-(2-Diethylamino-ethoxy)-phenylamino]-3-(3,5-dimethoxy-phenyl)-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2-yl]-3-ethyl-urea;

1-{3-(2-Chloro-3,5-dimethoxy-phenyl)-7-[4-(2-diethylamino-ethoxy)-phenylamino]-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2-yl}-3-ethyl-urea;

1-*tert*-Butyl-3-[7-[4-(2-diethylamino-ethoxy)-phenylamino]-3-(3,5-dimethoxy-phenyl)-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2-yl]-urea;

1-*tert*-Butyl-3-{3-(2-chloro-3,5-dimethoxy-phenyl)-7-[4-(2-diethylamino-ethoxy)-phenylamino]-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2-yl}-urea;
 1-*tert*-Butyl-3-[3-(3,5-dimethoxy-phenyl)-7-(pyridin-4-ylamino)-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2-yl]-urea;
 1-[3-(3,5-Dimethoxy-phenyl)-7-(pyridin-4-ylamino)-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2-yl]-3-ethyl-urea;
 1-*tert*-Butyl-3-[3-(2-chloro-3,5-dimethoxy-phenyl)-7-(pyridin-4-ylamino)-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2-yl]-urea;
 1-[3-(2-Chloro-3,5-dimethoxy-phenyl)-7-(pyridin-4-ylamino)-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2-yl]-3-ethyl-urea;
 1-[3-(2-Chloro-3,5-dimethoxy-phenyl)-7-(4-diethylamino-butylamino)-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2-yl]-3-ethyl-urea;
 3-Methyl-N-{7-[4-(5-methyl-hexahydro-pyrrolo[3,4-*c*]pyrrol-2-yl)-phenylamino]-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2-yl}-butyramide;
 1-{7-[4-(4-Acetyl-piperazin-1-yl)-phenylamino]-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2-yl}-3-isopropyl-urea; and
 1-*tert*-Butyl-3-[3-(2-chloro-3,5-dimethoxy-phenyl)-7-(4-diethylamino-butylamino)-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2-yl]-urea.

12. (currently amended) A compound of Claim 2 having the formula III




or a pharmaceutically acceptable salt thereof,

wherein:

R¹ is selected from the group consisting of H, (CH₂)_nAr, COR⁴, (CH₂)_nheteroaryl, (CH₂)_nheterocyclyl, C₁-C₁₀ alkyl, C₃-C₁₀ cycloalkyl, C₂-C₁₀ alkenyl, and C₂-C₁₀ alkynyl, wherein n is 0, 1, 2, or 3, and the (CH₂)_nAr, (CH₂)_nheteroaryl, alkyl, cycloalkyl, alkenyl, and alkynyl groups are optionally substituted by up to 5 groups selected from NR⁴R⁵, N⁺(O)R⁴R⁵, N⁺R⁴R⁵R⁶Y⁻, alkyl, phenyl, substituted phenyl, (CH₂)_nheteroaryl, hydroxy, alkoxy, phenoxy, thiol,

thioalkyl, halo, COR⁴, CO₂R⁴, CONR⁴R⁵, SO₂NR⁴R⁵, SO₃R⁴, PO₃R⁴, aldehyde, nitrile, nitro, heteroaryloxy, T(CH₂)_mQR⁴,



C(O)T(CH₂)_mQR⁴, NHC(O)T(CH₂)_mQR⁴, T(CH₂)_mC(O)NR⁴NR⁵, or T(CH₂)_mCO₂R⁴ wherein each m is independently 1-6, T is O, S, NR⁴, N⁺(O)R⁴, N⁺R⁴R⁶Y⁻, or CR⁴R⁵, and Q is O, S, NR⁵, N⁺(O)R⁵, or N⁺R⁵R⁶Y⁻;

R⁴ and R⁵ are each independently selected from the group consisting of hydrogen, C₁-C₆ alkyl, substituted alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, N(C₁-C₆alkyl)₁ or ₂, (CH₂)_nAr,

C₃-C₁₀ cycloalkyl, heterocyclyl, and heteroaryl, or R⁴ and R⁵ together with the nitrogen to which they are attached optionally form a ring having 3 to 7 carbon atoms and said ring optionally contains 1, 2, or 3 heteroatoms selected from the group consisting of nitrogen, substituted nitrogen, oxygen, and sulfur;

when R⁴ and R⁵ together with the nitrogen to which they are attached form a ring, the said ring is optionally substituted by 1 to 3 groups selected from OH, OR⁴, NR⁴R⁵, (CH₂)_mOR⁴,

(CH₂)_mNR⁴R⁵, T-(CH₂)_mQR⁴, CO-T-(CH₂)_mQR⁴, NH(CO)T(CH₂)_mQR⁴, T-(CH₂)_mCO₂R⁴, or T(CH₂)_mCONR⁴R⁵;

R⁶ is alkyl; and

Y is a halo counter-ion.

13. (original) A compound selected from:

1-[7-(4-Fluoro-phenylamino)-pyrimido[4,5-*d*]pyrimidin-2-yl]-3-methyl-urea;

1-Isopropyl-3-(7-phenylamino-pyrimido[4,5-*d*]pyrimidin-2-yl)-urea;

1-{7-[4-(3-Aminomethyl-pyrrolidin-1-yl)-phenylamino]-pyrimido[4,5-*d*]pyrimidin-2-yl}-3-isopropyl-urea;

1-Isopropyl-3-[7-(4-piperazin-1-yl-phenylamino)-pyrimido[4,5-*d*]pyrimidin-2-yl]-urea;

1-{7-[4-(4-Acetyl-piperazin-1-yl)-phenylamino]-pyrimido[4,5-*d*]pyrimidin-2-yl}-3-isopropyl-urea;


N-{7-[4-(3-Amino-pyrrolidin-1-yl)-phenylamino]-pyrimido[4,5-*d*]pyrimidin-2-yl}-3-methyl-butyramide;

N-[7-(4-Piperazin-1-yl-phenylamino)-pyrimido[4,5-*d*]pyrimidin-2-yl]-isobutyramide;

N-{7-[4-(4-Acetyl-piperazin-1-yl)-phenylamino]-pyrimido[4,5-*d*]pyrimidin-2-yl}-3-methyl-butyramide;

3-Methyl-N-[7-(pyridin-4-ylamino)-pyrimido[4,5-d]pyrimidin-2-yl]-butyramide;
 1-Isopropyl-3-[7-(pyridin-4-ylamino)-pyrimido[4,5-d]pyrimidin-2-yl]-urea; and
 N-[7-[4-(3-Aminomethyl-pyrrolidin-1-yl)-phenylamino]-pyrimido[4,5-d]pyrimidin-2-yl]-3-methyl-
 butyramide.

14. (currently amended) A compound of Formula IV Claim 1 wherein W is S, SO, or SO₂



IV

or a pharmaceutically acceptable salt thereof,

wherein:

the dotted line represents an optional double bond;


W is S, SO, or SO₂;

X is either O, S, or NR¹⁰;

R¹, R², and R¹⁰ are independently selected from the group consisting of H, (CH₂)_nAr, COR⁴, (CH₂)_nheteroaryl, (CH₂)_nheterocyclyl, C₁-C₁₀alkyl, C₃-C₁₀cycloalkyl, C₂-C₁₀alkenyl, and C₂-C₁₀alkynyl, wherein n is 0, 1, 2, or 3, and the (CH₂)_nAr, (CH₂)_nheteroaryl, alkyl, cycloalkyl, alkenyl, and alkynyl groups are optionally substituted by up to 5 groups selected from NR⁴R⁵, N⁺(O)R⁴R⁵, N⁺R⁴R⁵R⁶Y⁻, alkyl, phenyl, substituted phenyl, (CH₂)_nheteroaryl, hydroxy, alkoxy, phenoxy, thiol, thioalkyl, halo, COR⁴, CO₂R⁴, CONR⁴R⁵, SO₂NR⁴R⁵, SO₃R⁴, PO₃R⁴, aldehyde, nitrile, nitro,

_____OR⁵
_____|
heteroaryloxy, T(CH₂)_mQR⁴, T(CH₂)_mC-(CH₂)_mQR⁴,


_____|
_____H
C(O)T(CH₂)_mQR⁴, NHC(O)T(CH₂)_mQR⁴, T(CH₂)_mC(O)NR⁴NR⁵, or T(CH₂)_mCO₂R⁴ wherein
each m is independently 1-6, T is O, S, NR⁴, N⁺(O)R⁴, N⁺R⁴R⁶Y⁻, or CR⁴R⁵, and Q is O, S,
NR⁵, N⁺(O)R⁵, or N⁺R⁵R⁶Y⁻;



or a pharmaceutically acceptable salt thereof.


wherein:

R^1 and R^2 are independently selected from the group consisting of $(CH_2)_nAr$, COR^4 , $(CH_2)_n$ heteroaryl, $(CH_2)_n$ heterocyclyl, C_3-C_{10} cycloalkyl, C_2-C_{10} alkenyl, and C_2-C_{10} alkynyl, wherein n is 0, 1, 2, or 3, and the $(CH_2)_nAr$, $(CH_2)_n$ heteroaryl, alkyl, cycloalkyl, alkenyl, and alkynyl groups are optionally substituted by up to 5 groups selected from NR^4R^5 , $N^+(O)R^4R^5$, $N^+R^4R^5R^6Y^-$, alkyl, phenyl, substituted phenyl, $(CH_2)_n$ heteroaryl, hydroxy, alkoxy, phenoxy, thiol, thioalkyl, halo, COR^4 , CO_2R^4 , $CONR^4R^5$, $SO_2NR^4R^5$, SO_3R^4 , PO_3R^4 , aldehyde, nitrile, nitro, heteroaryloxy, $T(CH_2)_mQR^4$.



$C(O)T(CH_2)_mQR^4$, $NHC(O)T(CH_2)_mQR^4$, $T(CH_2)_mC(O)NR^4R^5$, or $T(CH_2)_mCO_2R^4$ wherein each m is independently 1-6, T is O, S, NR^4 , $N^+(O)R^4$, $N^+R^4R^6Y^-$, or CR^4R^5 , and Q is O, S, NR^5 , $N^+(O)R^5$, or $N^+R^5R^6Y^-$.

R^3 has the meanings of R^2 , wherein R^2 is as defined above, as well as OH, NR^4R^5 , $COOR^4$, OR^4 , $CONR^4R^5$, $SO_2NR^4R^5$, SO_3R^4 , PO_3R^4 , $T(CH_2)_mQR^4$.



wherein T and Q are as defined above;

R^4 and R^5 are each independently selected from the group consisting of hydrogen, C_1-C_6 alkyl, substituted alkyl, C_2-C_6 alkenyl, C_2-C_6 alkynyl, $N(C_1-C_6alkyl)_1$ or 2 , $(CH_2)_nAr$.

C₃-C₁₀ cycloalkyl, heterocyclyl, and heteroaryl, or R⁴ and R⁵ together with the nitrogen to which they are attached optionally form a ring having 3 to 7 carbon atoms and said ring optionally contains 1, 2, or 3 heteroatoms selected from the group consisting of nitrogen, substituted nitrogen, oxygen, and sulfur;

when R⁴ and R⁵ together with the nitrogen to which they are attached form a ring, the said ring is optionally substituted by 1 to 3 groups selected from OH, OR⁴, NR⁴R⁵, (CH₂)_mOR⁴,

(CH₂)_mNR⁴R⁵, T-(CH₂)_mQR⁴, CO-T-(CH₂)_mQR⁴, NH(CO)T(CH₂)_mQR⁴, T-(CH₂)_mCO₂R⁴, or T(CH₂)_mCONR⁴R⁵;

R⁶ is alkyl; and


Y is a halo counter-ion.

16. (original) A compound selected from:

1-Isopropyl-7-[4-(4-methylpiperazin-1-yl)phenylamino]-1*H*-pyrimido[4,5-*d*]pyrimidine-2,4-dione;
7-[4-(2-Diethylaminoethoxy)phenylamino]-1-isopropyl-1*H*-pyrimido[4,5-*d*]pyrimidine-2,4-dione;
7-(4-Diethylamino-butylamino)-3-(3,5-dimethoxy-phenyl)-1-ethyl-1*H*-pyrimido[4,5-*d*]pyrimidine-2,4-dione;
7-[4-(2-Diethylamino-ethoxy)-phenylamino]-3-(3,5-dimethoxy-phenyl)-1-ethyl-1*H*-pyrimido[4,5-*d*]pyrimidine-2,4-dione; and
7-(Pyridin-4-ylamino)-3-(3,5-dimethoxy-phenyl)-1-ethyl-1*H*-pyrimido[4,5-*d*]pyrimidine-2,4-dione.

Claims 17-25 (canceled).

26. (currently amended) A method of inhibiting a cyclin-dependent kinase comprising contacting the cyclin-dependent kinase with a compound of Formula VI



VI


or a pharmaceutically acceptable salt thereof, and the pharmaceutically acceptable salts thereof,
wherein:

the dotted line represents an optional double bond;

W is NH, S, SO, or SO₂;

X is either O, S, or NR¹⁰;


R¹, R², and R¹⁰ are independently selected from the group consisting of H, (CH₂)_nAr, COR⁴, (CH₂)_nheteroaryl, (CH₂)_nheterocyclyl, C₁-C₁₀ alkyl, C₃-C₁₀ cycloalkyl, C₂-C₁₀ alkenyl, and C₂-C₁₀ alkynyl, wherein n is 0, 1, 2, or 3, and the (CH₂)_nAr, (CH₂)_nheteroaryl, alkyl, cycloalkyl, alkenyl, and alkynyl groups are optionally substituted by up to 5 groups selected from NR⁴R⁵, N⁺(O)R⁴R⁵, N⁺R⁴R⁵R⁶Y⁻, alkyl, phenyl, substituted phenyl, (CH₂)_nheteroaryl, hydroxy, alkoxy, phenoxy, thiol, thioalkyl, halo, COR⁴, CO₂R⁴, CONR⁴R⁵, SO₂NR⁴R⁵, SO₃R⁴, PO₃R⁴, aldehyde, nitrile, nitro,



C(O)T(CH₂)_mQR⁴, NHC(O)T(CH₂)_mQR⁴, T(CH₂)_mC(O)NR⁴NR⁵, or T(CH₂)_mCO₂R⁴ wherein each m is independently 1-6, T is O, S, NR⁴, N⁺(O)R⁴, N⁺R⁴R⁶Y⁻, or CR⁴R⁵, and Q is O, S, NR⁵, N⁺(O)R⁵, or N⁺R⁵R⁶Y⁻;

when the dotted line is present, R³ is absent;

otherwise R³ has the meanings of R², wherein R² is as defined above, as well as OH, NR⁴R⁵, COOR⁴, OR⁴, CONR⁴R⁵, SO₂NR⁴R⁵, SO₃R⁴, PO₃R⁴,



wherein T and Q are as defined above;

R⁴ and R⁵ are each independently selected from the group consisting of hydrogen, C₁-C₆ alkyl, substituted alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, N(C₁-C₆alkyl)_{1 or 2}, (CH₂)_nAr,

C₃-C₁₀ cycloalkyl, heterocyclyl, and heteroaryl, or R⁴ and R⁵ together with the nitrogen to which they are attached optionally form a ring having 3 to 7 carbon atoms and said ring optionally contains 1, 2, or 3 heteroatoms selected from the group consisting of nitrogen, substituted nitrogen, oxygen, and sulfur;


when R⁴ and R⁵ together with the nitrogen to which they are attached form a ring, the said ring is optionally substituted by 1 to 3 groups selected from OH, OR⁴, NR⁴R⁵, (CH₂)_mOR⁴,

$(\text{CH}_2)_m\text{NR}^4\text{R}^5$, $\text{T}-(\text{CH}_2)_m\text{QR}^4$, $\text{CO-T}-(\text{CH}_2)_m\text{QR}^4$, $\text{NH(CO)T}-(\text{CH}_2)_m\text{QR}^4$, $\text{T}-(\text{CH}_2)_m\text{CO}_2\text{R}^4$, or $\text{T}-(\text{CH}_2)_m\text{CONR}^4\text{R}^5$;

R^6 is alkyl;

R^8 and R^9 independently are H, $\text{C}_1\text{-C}_3$ alkyl, NR^4R^5 , $\text{N}^+(\text{O})\text{R}^4\text{R}^5$, $\text{N}^+\text{R}^4\text{R}^5\text{R}^6\text{Y}^-$, hydroxy, alkoxy, thiol, thioalkyl, halo, COR^4 , CO_2R^4 , CONR^4R^5 , $\text{SO}_2\text{NR}^4\text{R}^5$, SO_3R^4 , PO_3R^4 , CHO, CN, or NO_2 ;

when the dotted line is absent, R^9 is additionally oxo,




Y is a halo counter-ion.

27. (currently amended) [[A]] The method of Claim 26 wherein said cyclin-dependent kinase is cdc2.

28. (currently amended) [[A]] The method of Claim 26 wherein said cyclin-dependent kinase is cdk2.

29. (currently amended) [[A]] The method of Claim 26 wherein said cyclin-dependent kinase is cdk4 or cdk6.

30. (currently amended) A method of inhibiting a growth factor-mediated tyrosine kinase comprising contacting said growth factor-mediated kinase with a compound of Formula VI



VI


or a pharmaceutically acceptable salt thereof, and the pharmaceutically acceptable salts thereof,
wherein:

the dotted line represents an optional double bond;

W is NH, S, SO, or SO_2 ;

X is either O, S, or NR¹⁰;


R¹, R², and R¹⁰ are independently selected from the group consisting of H, (CH₂)_nAr, COR⁴, (CH₂)_nheteroaryl, (CH₂)_nheterocyclyl, C₁-C₁₀ alkyl, C₃-C₁₀ cycloalkyl, C₂-C₁₀ alkenyl, and C₂-C₁₀ alkynyl, wherein n is 0, 1, 2, or 3, and the (CH₂)_nAr, (CH₂)_nheteroaryl, alkyl, cycloalkyl, alkenyl, and alkynyl groups are optionally substituted by up to 5 groups selected from NR⁴R⁵, N⁺(O)R⁴R⁵, N⁺R⁴R⁵R⁶Y⁻, alkyl, phenyl, substituted phenyl, (CH₂)_nheteroaryl, hydroxy, alkoxy, phenoxy, thiol, thioalkyl, halo, COR⁴, CO₂R⁴, CONR⁴R⁵, SO₂NR⁴R⁵, SO₃R⁴, PO₃R⁴, aldehyde, nitrile, nitro,



C(O)T(CH₂)_mQR⁴, NHC(O)T(CH₂)_mQR⁴, T(CH₂)_mC(O)NR⁴NR⁵, or T(CH₂)_mCO₂R⁴ wherein each m is independently 1-6, T is O, S, NR⁴, N⁺(O)R⁴, N⁺R⁴R⁶Y⁻, or CR⁴R⁵, and Q is O, S, NR⁵, N⁺(O)R⁵, or N⁺R⁵R⁶Y⁻;

when the dotted line is present, R³ is absent;

otherwise R³ has the meanings of R², wherein R² is as defined above, as well as OH, NR⁴R⁵, COOR⁴, OR⁴, CONR⁴R⁵, SO₂NR⁴R⁵, SO₃R⁴, PO₃R⁴,



wherein T and Q are as defined above;

R⁴ and R⁵ are each independently selected from the group consisting of hydrogen, C₁-C₆ alkyl, substituted alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, N(C₁-C₆alkyl)₁ or 2, (CH₂)_nAr,

C₃-C₁₀ cycloalkyl, heterocyclyl, and heteroaryl, or R⁴ and R⁵ together with the nitrogen to which they are attached optionally form a ring having 3 to 7 carbon atoms and said ring optionally contains 1, 2, or 3 heteroatoms selected from the group consisting of nitrogen, substituted nitrogen, oxygen, and sulfur;


when R⁴ and R⁵ together with the nitrogen to which they are attached form a ring, the said ring is optionally substituted by 1 to 3 groups selected from OH, OR⁴, NR⁴R⁵, (CH₂)_mOR⁴,

$(\text{CH}_2)_m\text{NR}^4\text{R}^5$, $\text{T}-(\text{CH}_2)_m\text{QR}^4$, $\text{CO-T}-(\text{CH}_2)_m\text{QR}^4$, $\text{NH(CO)T}-(\text{CH}_2)_m\text{QR}^4$, $\text{T}-(\text{CH}_2)_m\text{CO}_2\text{R}^4$, or $\text{T}-(\text{CH}_2)_m\text{CONR}^4\text{R}^5$;

R^6 is alkyl;

R^8 and R^9 independently are H, $\text{C}_1\text{-C}_3$ alkyl, NR^4R^5 , $\text{N}^+(\text{O})\text{R}^4\text{R}^5$, $\text{N}^+\text{R}^4\text{R}^5\text{R}^6\text{Y}^-$, hydroxy, alkoxy, thiol, thioalkyl, halo, COR^4 , CO_2R^4 , CONR^4R^5 , $\text{SO}_2\text{NR}^4\text{R}^5$, SO_3R^4 , PO_3R^4 , CHO, CN, or NO_2 ;

when the dotted line is absent, R^9 is additionally oxo,




Y is a halo counter-ion.

31. (currently amended) [[A]] The method of Claim 30 wherein said growth factor-mediated tyrosine kinase is platelet derived growth factor (PDGF).

32. (currently amended) [[A]] The method of Claim 30 wherein said growth factor-mediated tyrosine kinase is fibroblast growth factor (FGF).

33. (currently amended) [[A]] The method of Claim 30 wherein said growth factor-mediated tyrosine kinase is vascular endothelial growth factor (VEGF).

34. (currently amended) A method of inhibiting a non-receptor tyrosine kinase comprising contacting said non-receptor tyrosine kinase with a compound of Formula VI



VI


or a pharmaceutically acceptable salt thereof, and the pharmaceutically acceptable salts thereof,
wherein:

the dotted line represents an optional double bond;

W is NH, S, SO, or SO_2 ;

X is either O, S, or NR¹⁰;


R¹, R², and R¹⁰ are independently selected from the group consisting of H, (CH₂)_nAr, COR⁴, (CH₂)_nheteroaryl, (CH₂)_nheterocyclyl, C₁-C₁₀ alkyl, C₃-C₁₀ cycloalkyl, C₂-C₁₀ alkenyl, and C₂-C₁₀ alkynyl, wherein n is 0, 1, 2, or 3, and the (CH₂)_nAr, (CH₂)_nheteroaryl, alkyl, cycloalkyl, alkenyl, and alkynyl groups are optionally substituted by up to 5 groups selected from NR⁴R⁵, N⁺(O)R⁴R⁵, N⁺R⁴R⁵R⁶Y⁻, alkyl, phenyl, substituted phenyl, (CH₂)_nheteroaryl, hydroxy, alkoxy, phenoxy, thiol, thioalkyl, halo, COR⁴, CO₂R⁴, CONR⁴R⁵, SO₂NR⁴R⁵, SO₃R⁴, PO₃R⁴, aldehyde, nitrile, nitro,



C(O)T(CH₂)_mQR⁴, NHC(O)T(CH₂)_mQR⁴, T(CH₂)_mC(O)NR⁴NR⁵, or T(CH₂)_mCO₂R⁴ wherein each m is independently 1-6, T is O, S, NR⁴, N⁺(O)R⁴, N⁺R⁴R⁶Y⁻, or CR⁴R⁵, and Q is O, S, NR⁵, N⁺(O)R⁵, or N⁺R⁵R⁶Y⁻;

when the dotted line is present, R³ is absent;

otherwise R³ has the meanings of R², wherein R² is as defined above, as well as OH, NR⁴R⁵, COOR⁴, OR⁴, CONR⁴R⁵, SO₂NR⁴R⁵, SO₃R⁴, PO₃R⁴,



wherein T and Q are as defined above;

R⁴ and R⁵ are each independently selected from the group consisting of hydrogen, C₁-C₆ alkyl, substituted alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, N(C₁-C₆alkyl)₁ or 2, (CH₂)_nAr,

C₃-C₁₀ cycloalkyl, heterocyclyl, and heteroaryl, or R⁴ and R⁵ together with the nitrogen to which they are attached optionally form a ring having 3 to 7 carbon atoms and said ring optionally contains 1, 2, or 3 heteroatoms selected from the group consisting of nitrogen, substituted nitrogen, oxygen, and sulfur;

when R⁴ and R⁵ together with the nitrogen to which they are attached form a ring, the said ring is optionally substituted by 1 to 3 groups selected from OH, OR⁴, NR⁴R⁵, (CH₂)_mOR⁴,

$(\text{CH}_2)_m\text{NR}^4\text{R}^5$, $\text{T}-(\text{CH}_2)_m\text{QR}^4$, $\text{CO-T}-(\text{CH}_2)_m\text{QR}^4$, $\text{NH(CO)T}-(\text{CH}_2)_m\text{QR}^4$, $\text{T}-(\text{CH}_2)_m\text{CO}_2\text{R}^4$, or $\text{T}-(\text{CH}_2)_m\text{CONR}^4\text{R}^5$;

R^6 is alkyl;

R^8 and R^9 independently are H, $\text{C}_1\text{-C}_3$ alkyl, NR^4R^5 , $\text{N}^+(\text{O})\text{R}^4\text{R}^5$, $\text{N}^+\text{R}^4\text{R}^5\text{R}^6\text{Y}^-$, hydroxy, alkoxy, thiol, thioalkyl, halo, COR^4 , CO_2R^4 , CONR^4R^5 , $\text{SO}_2\text{NR}^4\text{R}^5$, SO_3R^4 , PO_3R^4 , CHO, CN, or NO_2 ;


when the dotted line is absent, R^9 is additionally oxo,

$\begin{array}{ccccc} \text{S} & \text{NH} & \text{NR}^4 & \text{NOH} & \text{NOR}^4 \\ || & || & || & || & || \\ \text{C-R}^4 & \text{CH} & \text{CH} & \text{CH} & \text{C} \end{array}$
 and

Y is a halo counter-ion.

35. (currently amended) [[A]] The method of Claim 3[[3]]4 wherein said non-receptor tyrosine kinase is selected from a transforming gene of the Rous sarcoma retrovirus (Src) family.

36. (currently amended) A method of inhibiting a serine kinase in a mammal comprising administering a serine kinase inhibiting among of a compound of ~~Claim 4~~ Formula VI



VI

or a pharmaceutically acceptable salt thereof,

wherein:


the dotted line represents an optional double bond;

W is NH, S, SO, or SO_2 ;

X is either O, S, or NR^{10} .

R^1 , R^2 , and R^{10} are independently selected from the group consisting of H, $(\text{CH}_2)_n\text{Ar}$, COR^4 , $(\text{CH}_2)_n\text{heteroaryl}$, $(\text{CH}_2)_n\text{heterocyclyl}$, $\text{C}_1\text{-C}_{10}\text{alkyl}$, $\text{C}_3\text{-C}_{10}\text{cycloalkyl}$, $\text{C}_2\text{-C}_{10}\text{alkenyl}$, and


C₂-C₁₀ alkynyl, wherein n is 0, 1, 2, or 3, and the (CH₂)_nAr, (CH₂)_nheteroaryl, alkyl, cycloalkyl, alkenyl, and alkynyl groups are optionally substituted by up to 5 groups selected from NR⁴R⁵, N⁺(O)R⁴R⁵, N⁺R⁴R⁵R⁶Y⁻, alkyl, phenyl, substituted phenyl, (CH₂)_nheteroaryl, hydroxy, alkoxy, phenoxy, thiol, thioalkyl, halo, COR⁴, CO₂R⁴, CONR⁴R⁵, SO₂NR⁴R⁵, SO₃R⁴, PO₃R⁴, aldehyde, nitrile, nitro,



C(O)T(CH₂)_mQR⁴, NHC(O)T(CH₂)_mQR⁴, T(CH₂)_mC(O)NR⁴NR⁵, or T(CH₂)_mCO₂R⁴ wherein each m is independently 1-6, T is O, S, NR⁴, N⁺(O)R⁴, N⁺R⁴R⁵Y⁻, or CR⁴R⁵, and Q is O, S, NR⁵, N⁺(O)R⁵, or N⁺R⁵R⁶Y⁻;

when the dotted line is present, R³ is absent;

otherwise R³ has the meanings of R², wherein R² is as defined above, as well as OH, NR⁴R⁵, COOR⁴, OR⁴, CONR⁴R⁵, SO₂NR⁴R⁵, SO₃R⁴, PO₃R⁴,



wherein T and Q are as defined above;

R⁴ and R⁵ are each independently selected from the group consisting of hydrogen, C₁-C₆ alkyl, substituted alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, N(C₁-C₆alkyl)_{1 or 2}, (CH₂)_nAr,

C₃-C₁₀ cycloalkyl, heterocyclyl, and heteroaryl, or R⁴ and R⁵ together with the nitrogen to which they are attached optionally form a ring having 3 to 7 carbon atoms and said ring optionally contains 1, 2, or 3 heteroatoms selected from the group consisting of nitrogen, substituted nitrogen, oxygen, and sulfur;


when R⁴ and R⁵ together with the nitrogen to which they are attached form a ring, the said ring is optionally substituted by 1 to 3 groups selected from OH, OR⁴, NR⁴R⁵, (CH₂)_mOR⁴,

(CH₂)_mNR⁴R⁵, T-(CH₂)_mQR⁴, CO-T-(CH₂)_mQR⁴, NH(CO)T(CH₂)_mQR⁴, T-(CH₂)_mCO₂R⁴, or T(CH₂)_mCONR⁴R⁵;

R⁶ is alkyl;

R^8 and R^9 independently are H, C_1-C_3 alkyl, NR^4R^5 , $N^+(O)R^4R^5$, $N^+R^4R^5R^6Y^-$, hydroxy, alkoxy, thiol, thioalkyl, halo, COR^4 , CO_2R^4 , $CONR^4R^5$, $SO_2NR^4R^5$, SO_3R^4 , PO_3R^4 , CHO, CN, or NO_2 ;

when the dotted line is absent, R^9 is additionally oxo,



Y is a halo counter-ion.

Claims 37-41 (canceled).


42. (original) A compound selected from:

- 7-[3-(Carboxy)-phenylamino]-3-(2,6-dichloro-phenyl)-1-methyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
- 7-[3-(N-Dimethylaminopropyl-carboxamide)-phenylamino]-3-(2,6-dichloro-phenyl)-1-methyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
- 7-[3-(N-Dimethylaminopropyl-carboxamide)-phenylamino]-3-(2,6-dichloro-3-hydroxy-phenyl)-1-methyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
- 7-[3-(Carboxy)-phenylamino]-3-(2,6-dichloro-3-hydroxy-phenyl)-1-methyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
- 3-(2,6-Dichloro-phenyl)-7-[4-(2-ethylamino-ethoxy)-phenylamino]-1-methyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
- 3-(2,6-Dichloro-3-hydroxy-phenyl)-7-[4-(2-ethylamino-ethoxy)-phenylamino]-1-methyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
- 7-[4-(Carboxamide)-phenylamino]-3-(2,6-dichloro-phenyl)-1-methyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
- 7-[4-(Carboxamide)-phenylamino]-3-(2,6-dichloro-3-hydroxy-phenyl)-1-methyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
- 3-(2,6-Dichloro-phenyl)-7-(3-hydroxymethyl-phenylamino)-1-methyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
- 3-(2,6-Dichloro-phenyl)-7-(4-morpholin-4-yl-phenylamino)-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
- 3-(2,6-Dichloro-3-hydroxy-phenyl)-1-methyl-7-(4-morpholin-4-yl-phenylamino)-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
- 3-(2,6-Dichloro-3-hydroxy-phenyl)-7-(3-hydroxymethyl-phenylamino)-1-methyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;

7-[4-(3-Carboxypropyl)-phenylamino]-3-(2,6-dichloro-phenyl)-1-methyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
 7-[4-(3-Carboxypropyl)-phenylamino]-3-(2,6-dichloro-3-hydroxy-phenyl)-1-methyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one;
 3-(2,6-Dichloro-phenyl)-7-[4-(formyl-phenylamino)- 1-methyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one; and
 3-(2,6-Dichloro-3-hydroxy-phenyl)-7-[4-(formyl-phenylamino)- 1-methyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidin-2(1*H*)-one.

Claim 43 (canceled).

44. (original) A compound of the formula



wherein:

R¹ is C₁-C₁₀ alkyl or (CH₂)_nAr;


R² is H, C₁-C₁₀ alkyl, or (CH₂)_nAr; and

R³ is Ar,

wherein n is 0, 1, 2 or 3;

Ar is phenyl or phenyl substituted with one or two groups selected from halo, alkyl, or substituted alkyl; or a pharmaceutically acceptable salt thereof.

45. (original) A compound of the formula



wherein R^2 is $(CH_2)_nAr$, n is 0, 1, 2 or 3, and Ar is phenyl or phenyl substituted by a 2-aminoethyl group,
or a pharmaceutically acceptable salt thereof.


46. (currently amended) A pharmaceutical formulation comprising [[a]] the compound or pharmaceutically acceptable salt thereof of Claim 3 in combination with a pharmaceutically acceptable carrier, diluent or excipient.

Claim 47 (canceled).

48. (currently amended) A pharmaceutical formulation comprising [[a]] the compound or pharmaceutically acceptable salt thereof of Claim 44 in combination with a pharmaceutically acceptable carrier, diluent or excipient.

49. (currently amended) A pharmaceutical formulation comprising [[a]] the compound or pharmaceutically acceptable salt thereof of Claim 45 in combination with a pharmaceutically acceptable carrier, diluent or excipient.

50. (original) A compound of the formula



or a pharmaceutically acceptable salt thereof.


51. (original) The compound 7-(4-diethylamino-butylamino)-3-(2-chloro-3,5-dimethoxy-phenyl)-1-ethyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidine-2(1*H*)-one.

52. (original) The compound 7-(4-diethylamino-butylamino)-3-(2-methyl-3,5-dimethoxy-phenyl)-1-ethyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidine-2(1*H*)-one.

53. (original) The compound 7-(4-diethylamino-butylamino)-3-(3,5-dimethoxy-phenyl)-1-cyclopentyl-3,4-dihydro-pyrimido[4,5-*d*]pyrimidine-2(1*H*)-one.

Claims 54-55 (canceled).

56. (currently amended) A compound of ~~Claim 55 having the formula~~ VII




VII

or a pharmaceutically acceptable salt thereof,


wherein:

R^1 and R^2 independently are hydrogen, C_1 - C_{10} alkyl, $(CH_2)_n$ Ar, $(CH_2)_n$ heteroaryl, C_3 - C_{10} cycloalkyl, or $(CH_2)_n$ heterocyclyl, wherein n is 0, 1, 2 or 3, and the $(CH_2)_n$ Ar, $(CH_2)_n$ heteroaryl, alkyl, cycloalkyl and $(CH_2)_n$ heterocyclyl groups are optionally substituted by up to 5 groups selected from NR^4R^5 , $N^+(O)R^4R^5$, $N^+R^4R^5R^6Y^-$, alkyl, phenyl, substituted phenyl, $(CH_2)_n$ heteroaryl, hydroxy, alkoxy, phenoxy, thiol, thioalkyl, halo, COR^4 , CO_2R^4 , $CONR^4R^5$, $SO_2NR^4R^5$, SO_3R^4 , PO_3R^4 , aldehyde, nitrile, nitro, heteroaryloxy, $T(CH_2)_mQR^4$,



or $T(CH_2)_mCO_2R^4$ wherein each m is independently 1-6, T is O, S, NR^4 , $N^+(O)R^4$, $N^+R^4R^6Y^-$, or CR^4R^5 , and Q is O, S, NR^5 , $N^+(O)R^5$, or $N^+R^5R^6Y^-$;

R^3 has the meanings of R^2 , wherein R^2 is as defined above, as well as OH, NR^4R^5 , $COOR^4$, OR^4 , $CONR^4R^5$, $SO_2NR^4R^5$, SO_3R^4 , PO_3R^4 ,



wherein T and Q are as defined above;

R⁴ and R⁵ are each independently selected from the group consisting of hydrogen, C₁-C₆ alkyl, substituted alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, N(C₁-C₆alkyl)₁ or 2, (CH₂)_nAr,

C₃-C₁₀ cycloalkyl, heterocyclyl, and heteroaryl, or R⁴ and R⁵ together with the nitrogen to which they are attached optionally form a ring having 3 to 7 carbon atoms and said ring optionally contains 1, 2, or 3 heteroatoms selected from the group consisting of nitrogen, substituted nitrogen, oxygen, and sulfur;

when R⁴ and R⁵ together with the nitrogen to which they are attached form a ring, the said ring is optionally substituted by 1 to 3 groups selected from OH, OR⁴, NR⁴R⁵, (CH₂)_mOR⁴,

(CH₂)_mNR⁴R⁵, T-(CH₂)_mQR⁴, CO-T-(CH₂)_mQR⁴, NH(CO)T(CH₂)_mQR⁴, T-(CH₂)_mCO₂R⁴, or T(CH₂)_mCONR⁴R⁵;

R⁶ is alkyl; and

Y is a halo counter-ion.

Claims 57-58 (canceled).

59. (original) A pharmaceutical formulation comprising a compound of Claim 56 in combination with a pharmaceutically acceptable carrier, diluent or excipient.

60. (original) A compound of Claim 56 wherein R¹ is alkyl, pyridyl, or phenyl, each optionally substituted with hydroxy, alkoxy, NR⁴R⁵, or T(CH₂)_mQR⁴.

Claims 61-66 (not entered).

Claim 67 (canceled).

Claims 68-74 (not entered).

Claim 75 (canceled).

Claims 76-80 (not entered).